

## **Kisspeptin**

Hormone signalling

### **Total Content**

10mg

### **Dilution**

2ml

### **Dosing Guidelines**

Libido / Hormone Support – 150–300mcg

Fertility Focus – 500mcg

### **Click Conversion**

150mcg – 3 clicks

200mcg – 4 clicks

250mcg – 5 clicks

300mcg – 6 clicks

500mcg – 10 clicks

### **Protocol**

Morning or Pre-Evening for libido support

2–3 x weekly / protocol dependant

4–8 weeks on

4 weeks off

### **Cycle Break Support Option**

PT 141

### **Pen Duration**

200 clicks total

150mcg (3 clicks): ~66 doses

200mcg (4 clicks): 50 doses

250mcg (5 clicks): 40 doses

300mcg (6 clicks): ~33 doses

500mcg (10 clicks): 20 doses

### **Disclaimer & Usage Guidance**

This protocol is provided for educational and informational purposes only. All compounds referenced are intended for research and therapeutic exploration and are not presented as medical advice, diagnosis, or treatment.

Dosing guidelines are illustrative and based on commonly reported practices. Individual requirements may vary significantly.

All peptides are highly individualised. Factors such as age, metabolism, mitochondrial health, lifestyle, and overall wellbeing play a significant role in determining response, benefits, and outcomes.

Cycling peptides is commonly used as a precautionary measure, although requirements may vary depending on the individual and protocol.

### **General Administration Guidance:**

Morning injections are typically performed in a fasted state; wait approximately 30 minutes before eating

Evening injections are best administered 2 hours after the last meal to optimise absorption

Any use of these compounds should be undertaken with appropriate medical oversight where applicable. The author

assumes no responsibility or liability for any outcomes resulting from the use or misuse of the information provided.